

*Citation for published version:*

Schneider, F, Parsons, S, Clift, S, Stolte, A & McManus, M 2018, 'Collected marine litter - A growing waste challenge', *Marine Pollution Bulletin*, vol. 128, pp. 162-174. <https://doi.org/10.1016/j.marpolbul.2018.01.011>

*DOI:*

[10.1016/j.marpolbul.2018.01.011](https://doi.org/10.1016/j.marpolbul.2018.01.011)

*Publication date:*

2018

*Document Version*

Peer reviewed version

[Link to publication](#)

*Publisher Rights*

CC BY-NC-ND

**University of Bath**

**Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Collected marine litter – A growing waste challenge

Falk Schneider<sup>1</sup>, Sophie Parsons<sup>1</sup>, Sally Clift<sup>1</sup>, Andrea Stolte<sup>2</sup>, Marcelle C. McManus<sup>1</sup>

<sup>1</sup>University of Bath, Department of Mechanical Engineering, Claverton Down, BA2 7AY, United Kingdom

<sup>2</sup>WWF Germany, WWF-Ostseebüro, Knieperwall 1, 18439 Stralsund, Germany

**Corresponding author:** Falk Schneider ([F.Schneider@bath.ac.uk](mailto:F.Schneider@bath.ac.uk))

---

### Abstract.

Marine litter, in particular plastic debris, poses a serious threat to marine life, human health and the economy. In order to reduce its impact, marine litter collections such as beach clean-ups are frequently conducted. This paper presents a systematic review of temporal developments, geographical distribution, quantities and waste treatment pathways of collected marine litter. Results from over 130 studies and projects highlight the world-wide increase in collection efforts. Many of these are in wealthy countries that do not primarily contribute to the problem. Over 250 thousand tonnes, have already been removed, but there is little or no information available regarding how this waste is treated or used post collection. This paper highlights the need for a whole-system quantitative assessment for the collection and waste treatment of marine litter, and identifies the challenges associated with utilising this waste in the future.

**Keywords** marine litter; life-cycle; systems perspective; collection; waste treatment

---

### 1. Introduction

Possibly the biggest landfill of our planet is the ocean. It accumulates various types of waste, called marine litter. This includes metals, glass, ceramics, textiles, paper and timber. Yet, the largest and probably most harmful fraction of marine litter is plastic debris. It is estimated that every year between 4.8 and 12.7 million tonnes of plastic waste enter the ocean from land-based sources alone (Jambeck et al., 2015). The high input is linked to plastics durability and lightweight which allows it unlike other types of marine litter to be easily transported via wind, waste water and rivers when it has been littered or inadequately disposed (Li, Tse and Fok, 2016). This is not only true for macroplastics but also for microplastics from e.g. car tire wear, clothes washing (Browne et al., 2011) and cosmetic products (Napper et al., 2015; Sherrington et al., 2016). Marine litter also originates from sea-based activities for example when fishing gear or cargo is lost, abandoned or discarded despite international legislation that prohibits the dumping of waste at sea (MARPOL Annex V, 1988; Jones, 1995).

Inside the ocean, marine litter is found in all seven ocean compartments: biota, coastlines, at the sea surface and the seafloor as well as in sediments, sea ice and the water column (Law, 2017; Figure 1). The contact with biota has been reported in 44 thousand reported cases of entanglement and ingestion affecting up to 1400 marine species worldwide (Gall and Thompson, 2015; Tekman et al., 2017a). Marine litter can also lead to a decline in tourism when washed ashore on beaches (McIlgorm, Campbell and Rule 2011; Keswani et al., 2016), destroy corals at the seafloor (Kühn, Bravo Rebolledo and van Franeker, 2015), spread invasive species (Kiessling, Gutow and Thiel, 2015) and cause accidents with ships (Cho, 2005). The plastic fraction of marine litter can be particularly harmful because it is able to travel over long time and distances before breaking down into smaller fragments (Andrady, 2011). Microplastics can then adsorb and release chemicals (Li, Tse and Fok,

2016), which become bioavailable after ingestion. This is also a risk for humans because these substances can accumulate along the biological food chain (Li, Tse and Fok, 2016).

On a political level, the United Nations Development Goals demand that the majority of countries worldwide “prevent and significantly reduce marine pollution” by 2025 (UN General Assembly, 2015). This may be achieved by implementing the increasingly popular framework of a closed-loop and resource efficient Circular Economy, as alternative to the established linear take-make dispose economy (Lieder and Rashid, 2016). The European Union has set this in motion in their “Action plan for the Circular Economy” implementing a waste hierarchy in which prevention, reuse, recycling and energy recovery are favoured over landfill in this respective order (European Commission, 2015). Whilst this may help to reduce the input of waste in the long run, in the meantime the collection of marine litter is essential to reduce its stock and impact in the ocean.

Reviews of marine litter collections are frequently undertaken (Li, Tse and Fok, 2016; Keswani et al., 2016; Iñiguez, Conesa, and Fullana, 2016; Browne et al., 2015; Barnes and Miller 2004; Derraik, 2002) but there is no published quantitative overview describing the output of the various collection efforts (Law, 2017). Also, such reviews traditionally adopt an upstream perspective that focuses on input reduction and the impact of marine litter in the ocean rather than downstream activities for the collected litter (Figure 1; but see Iñiguez, Conesa, and Fullana, 2016). Yet, the investigation of collection and further downstream pathways is crucial to understand the full environmental impact of marine litter which is a necessary input to policy-making in this area.

This study reviews current literature relating to the collection of marine litter, and uses the findings to support a case for greater systems thinking and a fuller life cycle perspective towards collection and downstream treatment activities. This would enable both collection and treatment approaches to meet the Circular Economy and Sustainable Development Goals. It identifies where current gaps lie which currently prevent us from developing a full systems approach, and specifies what further data would be needed to progress this.




Figure 1: Marine litter flow and perspective of this review

2. Materials and methods

2.1 Literature search

Based on the mode of collection the literature has been divided into two groups: scientific and non-scientific collections. Scientific collections are studies that describe the typically very structured collection process in

detail so that studies can be repeated and compared. For example, Alshawafi et al. (2017) marks several transects on a beach before starting the collection. Non-scientific collections on the other hand are result focused clean-up projects that do not commonly describe the collection process in detail.

#### *Scientific collections*

The literature search for scientific collections was based on peer-reviewed publications from the database Scopus using the title keywords “ocean”, “sea” or “marine” in conjunction with “plastic” or “debris”. After removing duplicates, 254 accessible articles published between January 2013 and April 2017 were evaluated in order to identify publications in which marine litter had been removed from the marine environment. Errata and reviews were excluded as were other publications when their methods section did not describe the marine litter collection protocol. This resulted in 103 publications.

#### *Non-scientific collections*

The search for non-scientific collections was based on references from the reviewed literature described above, the Global Ghost Gear Initiative (<http://www.ghostgear.org/>) and MARELITT (<http://www.marelitt.eu/>). The scientific publications and the Global Ghost Gear Initiative linked to individual removal projects worldwide whereas MARELITT had a focus on Europe specifically. In total 58 projects were identified for which a more detailed internet search was performed. For credibility, information was only taken from published documents and not from websites directly which gave 29 results.

## **2.2 Data extraction**

From the 103 scientific and 29 non-scientific collection projects data about the (1) removal period, (2) location, (3) ocean compartment, (4) removed quantity, (5) collection method and (6) waste treatment was extracted. Subsequently, this data was sorted in a descending order based on the quantity collected (Appendix 1 and 2). When projects described multiple collections, data about the removed quantity and collection method were taken separately for each represented ocean compartment. This lead to 114 and 32 rows of data entries for the scientific and non-scientific collection projects respectively (Appendix 1 and 2).

## **2.3 Assumptions and limitations**

Where a publication focused on derelict fishing gear and did not state otherwise it was assumed that the collection took place from the seafloor. Some publications give information in multiple formats, for example, by weight and by volume. To avoid double counting in these situations only one unit was extracted, giving priority in the order of: (1) weight, (2) item counts and (3) volume. For scientific collections from the coastline it was assumed that the collection was conducted manually if not otherwise specified.


This review is clearly limited to the approach and databases taken for the literature search. The approach focuses on marine litter collection but not on waste treatment pathways. Therefore such literature was excluded. The database Scopus has an emphasis on scientific studies but it has to be noted that the majority of marine litter is not reported in academic journals. Therefore other references, including the Global Ghost Gear Initiative and MARELITT were included, but this only provides a snapshot of the marine litter collections worldwide.

## **3. Results**

### **3.1 Historic development**

The starting date of the scientific and non-scientific collection projects indicate an increase of removal efforts over the last decade (Figure 2). This is possibly a response to the overall rise of awareness towards the problem following the long-term accumulation of plastic debris in the ocean (Petry and Benemann 2017). As studies

118 are necessarily published retrospectively, the recorded decline between 2013 and 2016 may well be explained  
 119 by the time needed to publish results for these recent time periods.



120 Figure 2: Development of marine litter collection efforts over time

121 Scientific collection projects are typically conducted over a limited and short period of time. Eight studies  
 122 reported data collected from between four and eight years, and nine studies covered data from ten years or  
 123 more. The remaining 71 studies for which data was available were conducted over a period of less than three  
 124 years, of which the majority were single collection efforts that were started and completed within the same  
 125 year.

126 Non-scientific collection projects are frequently designed for the long-term with no fixed end-date. Organisa-  
 127 tions typically publish the results of their collection efforts regularly, for example the International Coastal  
 128 Cleanup on an annual basis (International Coastal Cleanup and Ocean Conservancy 2016). Yet, there are also  
 129 clean-ups that are limited in time, for example after extreme weather events (Swanson et al., 2016) or when  
 130 new collection approaches are tested (WWF Poland, 2015).

131

132 **3.2 Geographical distribution**

133 Figure 3 positions the reviewed marine litter collections in a world map based on Jambeck et al. (2015) so that  
 134 the origin of plastic debris can be directly compared with the site of collection. Europe, North America and  
 135 Australia undertake major collection efforts but also Brazil and South Korea are strongly represented.

136 Scientific collection projects tend to focus on the Mediterranean Sea, Brazil, the Pacific Ocean and Australia  
 137 whereas non-scientific collection projects are typically conducted in the northern hemisphere. There is a nota-  
 138 ble gap between scientific and non-scientific efforts in South-America, Australia and Asia when South-Korea  
 139 is excluded. This may suggest that there is further potential for scientific knowledge to trigger clean-up projects  
 140 outside academia.

141 Russia, East and West Africa and the Western part of South America with the exception of Chile were not  
 142 identified as locations of marine litter collections (Figure 3). Yet, scientific and non-scientific collection pro-  
 143 jects are undertaken worldwide and therefore additional investigation is needed to determine whether those  
 144 areas have been studied before.

145




Figure 3: Geographical distribution of marine debris collections worldwide (map based on Jambeck et al., 2015), where collection locations were ascribed to regions or countries as whole

### 3.3 Systems perspective

A systems perspective looks at the interactions of single elements combining them to a whole picture which allows a better understanding of the context and implications of the system in its embedded environment. In this way the ocean compartments, quantities, collection methods and waste treatment pathways were joined (Figure 4).

#### *Quantities and ocean compartments*

Within the reviewed literature quantitative data was not provided for 42 of 146 collections and the 104 collections that provided quantitative data collected 259,822 tonnes, 1,121,734 items and 1,586 dm<sup>3</sup> of marine litter from the ocean (Figure 4).

Table 1: Number of scientific and non-scientific collections per ocean compartment

	Scientific collections		Non-scientific collections		Total collections	
	Number	Percentage	Number	Percentage	Number	Percentage
Coastlines	44	39%	10	31%	54	37%
Biota	32	28%	0	0%	32	22%
Sea surface	24	21%	3	9%	27	18%
Sea ice	0	0%	0	0%	0	0%
Seafloor	7	6%	19	59%	26	18%
Water column	1	1%	0	0%	1	1%
Sediments	6	5%	0	0%	6	4%
Total	114		32		146	




Figure 4: Flow of marine litter from the ocean to waste treatment



161 The 114 scientific collections that were described in the peer-reviewed publications covered all ocean com-  
162 partments apart from the sea ice (Figure 4; Appendix 1). Quantitative data on the collected amounts were not  
163 available for 37% of the scientific collections including the one study collecting marine debris from the water  
164 column. For the remaining ocean compartments quantitative data was provided in 85.7%, 72.3%, 66.7%, 59.4%  
165 and 54.2% of the cases for the seafloor, sediments, coastlines, biota and the sea surface respectively. Most  
166 marine litter was collected from coastlines, which was also the most popular destination for removal activities  
167 (Table 1; Figure 4; Appendix 1).

168 The 32 non-scientific collections which were conducted outside academia removed marine litter from three  
169 out of seven ocean compartments; coastlines, sea surface and seafloor. Quantitative data was available for all  
170 collections. The largest amount of marine litter in terms of weight, item count and volume was picked up at  
171 coastlines followed by the seafloor and sea surface respectively. Non-scientific seafloor collections were un-  
172 dertaken almost twice as often as non-scientific coastline collections (Table 1, Figure 4; Appendix 2). This is  
173 unexpected given the large amount of regular beach clean-ups for example in local communities involving  
174 schools or local authorities. However, the relatively low number of non-scientific collections from coastlines  
175 can be explained by the review approach chosen in this study where non-scientific collections derived from  
176 websites such as the global ghost gear initiative that focuses on seafloor collections.

### 177 *Collection methods*

178 The collection methods used to remove marine litter from the ocean were typically described or implicitly  
179 given for scientific collections but only in 56.3% of the non-scientific collections (Figure 4; Appendix 1 and  
180 2). Within the literature that contained information on collection methods, 16 ocean compartment specific  
181 ways of collection were identified (Figure 4; Appendix 1 and 2). The scientific collection methods typically  
182 focused on the removal of marine litter by hand or by smaller gear whereas non-scientific collections outside  
183 academia generally involved larger machines such as cranes.

184 The collection methods that were reported within both scientific and non-scientific collections were: (1) man-  
185 ual removal of marine litter from the coastline, (2) surface trawling and (3) use of a retention boom to capture  
186 floating marine litter at the sea surface as well as (4) bottom trawling and (5) diving to pick up marine litter  
187 from the seafloor (Figure 4; Appendix 1 and 2). For manual collections from the coastline typically gloves and  
188 plastic bags are used to protect hands and store the marine litter. A retention boom is a floating barrier that  
189 collects buoyant litter passively by making use of water currents. During surface and bottom trawling nets are  
190 towed behind a ship although with different mesh size so that mainly smaller items are collected from the sea  
191 surface and larger items from the seafloor possibly as by-catch from normal fishing activities. Also diving is  
192 used to collect litter from the seafloor although it is restricted to shallow waters.

193 Scientific collection methods include: (1) flotation, (2) freeing, (3) necropsy, (4) Niskin bottles, (5) manual  
194 collection of marine litter from the sea surface and (6) sediment (Figure 4; Appendix 1). Also, (7) diving, (8)  
195 remotely operated vehicles (ROVs) and (9) grabs were used (Figure 4; Appendix 1). Flotation separates buoy-  
196 ant, typically microplastics from beach sediment using sea water and sieves. Freeing describes the attempt to  
197 remove marine litter from entangled animals typically to save their life. Necropsy is the removal and content  
198 analysis of a dead animals' intestine tract which is frequently conducted to study the impact of plastic debris  
199 ingestion on marine life. Niskin bottles are used to take samples from the water column. At the sea surface  
200 manual collections were conducted from ships whereas sediment samples were taken by divers or from shallow  
201 waters directly. In greater depth grabs and ROVs can be used to collect sediments.

202 Within the non-scientific collections marine litter was removed from the seabed using a (1) crane excavation  
203 and by employing a (2) combination of ROVs and divers (Figure 4; Appendix 2). The crane operates from a  
204 ship which makes it mobile in the ocean. The benefit of using ROVs and divers together is that ROVs can  
205 detect marine litter and divers can assist retrieving them if needed.

### 206 *Waste treatment*

207 For the waste treatment of the collected marine litter almost no information is available. In fact, only 27.6% of  
208 the publications describing non-scientific collections mentioned the waste treatment of marine litter whereas



209 it was not mentioned in scientific collections at all. This is noteworthy, especially for larger removal activities  
210 that collected up to ten tonnes and more (Polasek, Bering et al., 2017) because the question about their final  
211 treatment remains unanswered.

212 The non-scientific collections that described the waste treatment of collected marine litter cover all available  
213 treatment options: reuse, recycle, energy recovery and landfill. Prevention of marine litter implies a reduction  
214 of waste input into the ocean. As the input of waste happens much earlier in the life cycle of marine litter  
215 compared to its treatment after collection, prevention is not applicable at this point (Figure 4).

216 In one study, derelict fishing gear was collected from the ocean, cleaned and subsequently reused for its former  
217 function (FUNDY NORTH Fisherman's Association, 2016). Recycling was also possible for other materials  
218 for example metals (National Fish and Wildlife Foundation, 2016) such as lead line (Northwest Straits Foun-  
219 dation, 2015) or polystyrene buoys (Iñiguez, Conesa, and Fullana, 2016). Yet, material recycling is challenging  
220 because collected marine litter is usually contaminated and requires several pre-treatment steps including the  
221 separation of sand from plastics and cleaning (Iñiguez, Conesa, and Fullana, 2016). Commonly encountered  
222 small fluctuating quantities of collected marine litter (Fishing for litter Scotland, 2015; 2017) as well as a low  
223 oil price pose further challenges to recycling because such conditions do not favour investments which are  
224 needed to set up a recycling plant. Apart from material recycling, marine litter was also turned into art (Ghost  
225 nets Australia, n.d.; Olive Ridley Project, 2017). Energy recovery from marine litter was possible in some  
226 cases (Swanson et al., 2016; Iñiguez, Conesa, and Fullana, 2016; National Fish and Wildlife Foundation, 2016)  
227 but not in others due to its elevated heating (Fishing for litter Scotland, 2015; 2017). Due to those complications  
228 some parts, if not all (Fishing for litter Scotland, 2015; 2017), of the collected litter was disposed in landfills  
229 (Figure 4; Appendix 2).

230

#### 231 4. Discussion

232 Marine litter collections are not new but there has been a notable rise in reported activity within the last ten  
233 years, with rising awareness and visibility of the plastic debris problem. Although collection efforts may in-  
234 crease, it does not necessarily reduce the overall stock of marine litter in the ocean because a growing and  
235 much larger amount of new waste, in particular plastics, is added to the marine environment every year  
236 (Jambeck et al., 2015). It needs both a drastic input reduction and increase of collection efforts to achieve the  
237 Sustainable Development Goal 14.1 which requires an input prevention and a significant stock decline by 2025  
238 (United Nations General Assembly, 2015). For performance measurements and estimates on when the accu-  
239 mulation of marine litter might begin to decline, data on the current stock, annual in- and output are needed.  
240 This review is a first step in collating such output data on a global scale but further investigation is obviously  
241 required.

242 The collection of marine litter is undertaken worldwide but it may be expected that regions suffering from  
243 marine litter are conducting more clean-up activities than other regions. Yet, from the top ten contributors of  
244 marine plastic debris: China, Indonesia, Philippines, Vietnam, Sri Lanka, Thailand, Egypt, Malaysia, Nigeria  
245 and Bangladesh (Jambeck et al., 2015) only Malaysia, China and the Philippines lead collection projects within  
246 the reviewed literature. A possible explanation for this is that contributing countries are not necessarily the  
247 countries who suffer from the pollution, although this seems very unlikely because waste does not just disap-  
248 pear after its disposal. It appears more plausible to link collection efforts to the wealth of a country because  
249 resources, a functional waste management infrastructure, education and awareness are needed to provide the  
250 basis for clean-up activities. It is also possible that collections exist but were not reported.

251 In current literature collected marine litter is quantified by weight, item counts or volume when values are  
252 given at all. This makes a direct comparison difficult. It would be beneficial to express the collected quantities  
253 as a single unit, especially for the large amounts that become relevant from a waste treatment perspective.  
254 Nevertheless, the results show that coastline collections yield by far the largest amounts, followed by the sea-  
255 floor and subsequently the remaining ocean compartments. The inclusion of additional data from local author-  
256 ities would be expected to support this observation, as coastlines are routinely cleaned. This also suggests, that  
257 the amount of collected marine litter analysed in this study is a low estimate. On the other hand it can be argued

that the data from non-scientific collections may be overestimated and inaccurate: They may also include stones, sand and water which do not fall under the definition of marine litter but have a relatively high weight compared to plastic debris. The distinction of marine litter is especially difficult for coastline collections where it is not clear whether the litter originated from the ocean or from land. Thus, the numbers should be taken with all due care.

Within the reviewed literature much of the marine litter was collected manually. This indicates ample room for process development to automate current collection methods and make them more efficient. Yet, an automation implies the use of machines that entail environmental impacts from upstream activities such as energy generation. Environmental impacts can also be more direct for example to the seafloor from dragging operations (Erftemeijer et al., 2012) and it needs careful consideration as to whether a collection realises more benefits than it does harm. In general collections that can be conducted as part of the normal human presence in the marine environment for example beach clean-ups or the removal of marine litter by-catch from fishing nets appear the most cost effective and environmentally friendly solutions.

Information on the material composition of the collected marine litter is important for waste treatment companies because it determines the required pre-treatment steps for each waste treatment approach. A limitation of this study is that representative information on the weight-based material composition could not be compiled, because the data was only available for few large-scale collections. In those, litter from the coastlines comprised of 60% plastic, 15% rope and netting, 13% foam, 10% other including rubber and 2% metal (Polasek et al., 2017) whereas litter from the seafloor contained 52% to 76% plastics including derelict fishing gear, 5% to 11% metal, 4% to 14% rubber, 3% to 18% timber, 2% to 5% textiles and up to 20% other (Fishing for litter Scotland, 2015; Lueiro, 2013; Naturschutzbund Deutschland, 2015). Despite large uncertainties this confirms that collected marine litter is a very diverse mix of materials that is especially difficult to separate, clean and recycle. However, as incineration and landfilling is not encouraged within the Circular Economy, information on alternative waste treatment options for marine litter is needed.

In understanding challenges related to marine litter management it is vital that a systems perspective is sought. Apart from ocean compartments, quantities, collection methods and waste treatment options which have been linked in this study other elements of the supply chain such as transportation and the harbour reception need to be included in this system. Thus, a quantitative environmental assessment of the system is needed to confirm that the best possible collection methods and subsequent treatment options are used. It also ensures that both the direct collection and processing route is optimised to be most environmentally beneficial, and that any indirect consequences arising from the utilisation of marine litter are also included. In order to do this, further data about material flows and process data are needed.

290

## 291 5. Conclusions

Continuous waste input to the ocean and its accumulation in the marine environment has severe impacts on marine life, humans and the economy. This includes accidents and harm caused by entanglement and ingestion, the spread of invasive species across the ocean, the extinction of corals and a decline in tourism. While research is still conducted to understand the full extent of its impacts, the literature also shows that initiatives have already started to respond to the problem.

The immediate short-term solution that was analysed in this study was the collection of marine litter from the ocean. In 103 scientific studies and 29 projects from outside academia more than 250 thousand tonnes, 1 million items and 1.5 m<sup>3</sup> of marine litter have been removed from all ocean compartments apart from the sea ice using at least 16 different collection methods. From those the manual removal of marine litter from the coastline contributed most to the overall collection. However, with all this effort and an increasing number of ongoing projects it is not possible so far to counteract the yearly input of plastic waste to the ocean. This implies that the amount of marine litter is still increasing despite ongoing collection effort. In order to meet the Sustainable Development Goals much more marine litter must be collected and it should be further investigated which collection method is the most suitable to do so.

306 Increased collection efforts also spark the interest in the waste treatment of collected marine litter. Yet, none  
307 of the 103 scientific studies mentioned such pathways after collection. However, within the non-scientific  
308 projects landfilling and energy recovery were described as common pathways but also recycling and reuse was  
309 possible for some parts of the collected litter. In light of the Circular Economy and its waste hierarchy further  
310 investigation on waste treatment options for marine litter is necessary to minimise current and future environ-  
311 mental impacts.

312 A full system quantitative assessment from the impact of litter collections on the marine environment all the  
313 way to reuse and recycling options is needed to help policy makers and waste treatment companies identify  
314 the collection and treatment pathways of marine litter that are most environmentally friendly and minimise  
315 undesired side effects. This requires data and further study of material flows as well as process data.

316

## 317 **Acknowledgements**

318 This work was supported by a PhD studentship from the University of Bath.

319

## 320 **References**

321 Abreo, N.A.S., Macusi, E. D., Blatchley, D.D. and Cuenca, G.C. (2016). Ingestion of Marine Plastic Debris by  
322 Green Turtle (*Chelonia mydas*) in Davao Gulf, Mindanao, Philippines. *Philippine Journal of Science*, 145(1),  
323 17-23.

324 Acampora, H., Lyashevskaya, O., Van Franeker, J. A., and O'Connor, I. (2016). The use of beached bird surveys  
325 for marine plastic litter monitoring in Ireland. *Mar Environ Res*, 120, 122-129. doi:10.1016/j.maren-  
326 vres.2016.08.002

327 Agustin, A. E., Merrifield, M. A., Potemra, J. T., and Morishige, C. (2015). Temporal variability of marine de-  
328bris deposition at Tern Island in the Northwestern Hawaiian Islands. *Mar Pollut Bull*, 101(1), 200-207.  
329 doi:10.1016/j.marpolbul.2015.10.076

330 Alshawafi, A., Analla, M., Alwashali, E., and Aksissou, M. (2017). Assessment of marine debris on the coastal  
331 wetland of Martil in the North-East of Morocco. *Mar Pollut Bull*, 117(1-2), 302-310. doi:10.1016/j.marpol-  
332 bul.2017.01.079

333 Anastasopoulou, A., Mytilineou, C., Smith, C. J., and Papadopoulou, K. N. (2013). Plastic debris ingested by  
334 deep-water fish of the Ionian Sea (Eastern Mediterranean). *Deep Sea Research Part I: Oceanographic Re-  
335 search Papers*, 74, 11-13. doi:10.1016/j.dsr.2012.12.008

336 Andrady, A. (2011). Microplastics in the marine environment. *Mar Pollut Bull*, 62(8), 1596-1605.  
337 doi:10.1016/j.marpolbul.2011.05.030

338 Arun Kumar, A., Sivakumar, Sai Rutwik Reddy, Y., Bhagya Raja, M.V., Nishanth, T. and Revanth, V. (2016).  
339 Preliminary study on marine debris pollution along Marina beach, Chennai, India. *Regional Studies in Ma-  
340 rine Science*, 5, 35-40. doi:10.1016/j.rsma.2016.01.002

341 Avery-Gomm, S., Provencher, J. F., Morgan, K. H., and Bertram, D. F. (2013). Plastic ingestion in marine-as-  
342 sociated bird species from the eastern North Pacific. *Mar Pollut Bull*, 72(1), 257-259. doi:10.1016/j.marpol-  
343 bul.2013.04.021

344 Barnes, D. K. A. and Milner, P. (2004). Drifting plastic and its consequences for sessile organism dispersal in  
345 the Atlantic Ocean. *Marine Biology*, 146(4), 815-825.

346 Battaglia, P., Pedà, C., Musolino, S., Esposito, V., Andaloro, F., and Romeo, T. (2015). Diet and first docu-  
347 mented data on plastic ingestion of *Trachinotus ovatus* L. 1758 (Pisces: Carangidae) from the Strait of Mes-  
348 sina (central Mediterranean Sea). *Italian Journal of Zoology*, 83(1), 121-129.  
349 doi:10.1080/11250003.2015.1114157

350 Bennett-Martin, P., Visaggi, C. C., and Hawthorne, T. L. (2015). Mapping marine debris across coastal com-  
351 munities in Belize: developing a baseline for understanding the distribution of litter on beaches using geo-  
352 graphic information systems. *Environ Monit Assess*, 188(10), 557. doi:10.1007/s10661-016-5544-4

353 Blaskovic, A., Fastelli, P., Cizmek, H., Guerranti, C., and Renzi, M. (2017). Plastic litter in sediments from the  
354 Croatian marine protected area of the natural park of Telascica bay (Adriatic Sea). *Mar Pollut Bull*, 114(1),  
355 583-586. doi:10.1016/j.marpolbul.2016.09.018

356 Bond, A., Provencher, J., Elliot, R., Ryan, P., Rowe, S., Jones, I., Robertson, G. and Wilhelm, S. (2013). Inges-  
357 tion of plastic marine debris by Common and Thick-billed Murres in the northwestern Atlantic from 1985 to  
358 2012. *Marine Pollution Bulletin*, 77(1-2), pp.192-195. doi:10.1016/j.marpolbul.2013.10.005

359 Browne, M. A., Chapman, M. G., Thompson, R. C., Amaral Zettler, L. A., Jambeck, J., and Mallos, N. J. (2015).  
360 Spatial and temporal patterns of stranded intertidal marine debris: is there a picture of global change? *Envi-  
361 ron Sci Technol*, 49(12), 7082-7094. doi:10.1021/es5060572

362 Browne, M., Crump, P., Niven, S., Teuten, E., Tonkin, A., Galloway, T. and Thompson, R. (2011). Accumula-  
363 tion of Microplastic on Shorelines Worldwide: Sources and Sinks. *Environmental Science & Technology*,  
364 45(21), pp.9175-9179.

365 Campani, T., Bains, M., Giannetti, M., Cancelli, F., Mancusi, C., Serena, F., Marsili, L., Casini, S. and Fossi, M.  
366 (2013). Presence of plastic debris in loggerhead turtle stranded along the Tuscany coasts of the Pelagos  
367 Sanctuary for Mediterranean Marine Mammals (Italy). *Marine Pollution Bulletin*, 74(1), pp.225-230.  
368 doi:10.1016/j.marpolbul.2013.06.053

369 Carson, H. S., Lamson, M. R., Nakashima, D., Toloumu, D., Hafner, J., Maximenko, N. and McDermid, K. J.  
370 (2013). Tracking the sources and sinks of local marine debris in Hawai'i. *Mar Environ Res*, 84, 76-83.  
371 doi:10.1016/j.marenvres.2012.12.002

372 Casale, P., Freggi, D., Paduano, V., and Oliverio, M. (2016). Biases and best approaches for assessing debris  
373 ingestion in sea turtles, with a case study in the Mediterranean. *Mar Pollut Bull*, 110(1), 238-249.  
374 doi:10.1016/j.marpolbul.2016.06.057

375 Cheung, P. K., Cheung, L. T., and Fok, L. (2016). Seasonal variation in the abundance of marine plastic debris  
376 in the estuary of a subtropical macro-scale drainage basin in South China. *Sci Total Environ*, 562, 658-665.  
377 doi:10.1016/j.scitotenv.2016.04.048

378 Cho, D. O. (2005). Challenges to Marine Debris Management in Korea. *Coastal Management*, 33(4), 389-  
379 409.

380 Correa-Herrera, T., Barletta, M., Lima, A. R. A., Jimenez-Segura, L. F., and Arango-Sanchez, L. B. (2017). Spa-  
381 tial distribution and seasonality of ichthyoplankton and anthropogenic debris in a river delta in the Carib-  
382 bean Sea. *J Fish Biol*, 90(4), 1356-1387. doi:10.1111/jfb.13243

383 Cozar, A., Echevarria, F., Gonzalez-Gordillo, J., Irigoien, X., Ubeda, B., Hernandez-Leon, S., Palma, A., Na-  
384 varro, S., Garcia-de-Lomas, J., Ruiz, A., Fernandez-de-Puelles, M. and Duarte, C. (2014). Plastic debris in the  
385 open ocean. *Proceedings of the National Academy of Sciences*, 111(28), pp.10239-10244.  
386 doi:10.1073/pnas.1314705111

387 Cózar, A., Sanz-Martín, M., Martí, E., González-Gordillo, J., Ubeda, B., Gálvez, J., Irigoien, X. and Duarte, C.  
388 (2015). Plastic Accumulation in the Mediterranean Sea. PLOS ONE, 10(4), p.e0121762. doi:10.1371/jour-  
389 nal.pone.0121762

390 Cressey, D. (2016). The plastic ocean scientists know that there is a colossal amount of plastic in the oceans.  
391 But they don't know where it all is, what it looks like or what damage it does. Nature (536), 263-265.

392 Da Silva, M. L., de Araujo, F. V., Castro, R. O., & Sales, A. S. (2015a). Spatial-temporal analysis of marine de-  
393 bris on beaches of Niteroi, RJ, Brazil: Itaipu and Itacoatiara. Mar Pollut Bull, 92(1-2), 233-236.  
394 doi:10.1016/j.marpolbul.2014.12.036

395 Da Silva Mendes, S., de Carvalho, R. H., de Faria, A. F., and de Sousa, B. M. (2015b). Marine debris ingestion  
396 by Chelonia mydas (Testudines: Cheloniidae) on the Brazilian coast. Mar Pollut Bull, 92(1-2), 8-10.  
397 doi:10.1016/j.marpolbul.2015.01.010

398 Da Silva, M. L., Sales, A. S., Martins, S., Castro, R. O. and Araujo, F. V. (2016). The influence of the intensity  
399 of use, rainfall and location in the amount of marine debris in four beaches in Niteroi, Brazil: Sossego, Cam-  
400 boinhas, Charitas and Flechas. Mar Pollut Bull, 113(1-2), 36-39. doi:10.1016/j.marpolbul.2016.10.061

401 Davis, W., 3rd, and Murphy, A. G. (2015). Plastic in surface waters of the Inside Passage and beaches of the  
402 Salish Sea in Washington State. Mar Pollut Bull, 97(1-2), 169-177. doi:10.1016/j.marpolbul.2015.06.019

403 De Carvalho, R. H., Lacerda, P. D., da Silva Mendes, S., Barbosa, B. C., Paschoalini, M., Prezoto, F., and de  
404 Sousa, B. M. (2015). Marine debris ingestion by sea turtles (Testudines) on the Brazilian coast: an underesti-  
405 mated threat? Mar Pollut Bull, 101(2), 746-749. doi:10.1016/j.marpolbul.2015.10.002

406 De Lucia, G., Caliani, I., Marra, S., Camedda, A., Coppa, S., Alcaro, L., Campani, T., Giannetti, M., Coppola, D.,  
407 Cicero, A., Panti, C., Baini, M., Guerranti, C., Marsili, L., Massaro, G., Fossi, M. and Matiddi, M. (2014).  
408 Amount and distribution of neustonic micro-plastic off the western Sardinian coast (Central-Western Medi-  
409 terranean Sea). Marine Environmental Research, 100, pp.10-16. doi:10.1016/j.marenvres.2014.03.017

410 De Scisciolo, T., Mijts, E. N., Becker, T. and Eppinga, M. B. (2016). Beach debris on Aruba, Southern Carib-  
411 bean: Attribution to local land-based and distal marine-based sources. Mar Pollut Bull, 106(1-2), 49-57.  
412 doi:10.1016/j.marpolbul.2016.03.039

413 De Tender, C. A., Devriese, L. I., Haegeman, A., Maes, S., Ruttink, T., and Dawyndt, P. (2015). Bacterial Com-  
414 munity Profiling of Plastic Litter in the Belgian Part of the North Sea. Environ Sci Technol, 49(16), 9629-  
415 9638. doi:10.1021/acs.est.5b01093

416 Debrot, A. O., Meesters, H. W., Bron, P. S., and de Leon, R. (2013). Marine debris in mangroves and on the  
417 seabed: largely-neglected litter problems. Mar Pollut Bull, 72(1), 1. doi:10.1016/j.marpolbul.2013.03.023

418 Derraik, J. (2002). The pollution of the marine environment by plastic debris: a review. Marine Pollution  
419 Bulletin, 44(9), pp.842-852.

420 Di Benedetto, A. P., and Awabdi, D. R. (2014). How marine debris ingestion differs among megafauna spe-  
421 cies in a tropical coastal area. Mar Pollut Bull, 88(1-2), 86-90. doi:10.1016/j.marpolbul.2014.09.020

422 Di Benedetto, A. P., and Ramos, R. M. (2014). Marine debris ingestion by coastal dolphins: what drives dif-  
423 ferences between sympatric species? Mar Pollut Bull, 83(1), 298-301. doi:10.1016/j.marpolbul.2014.03.057

424 Duhec, A. V., Jeanne, R. F., Maximenko, N. and Hafner, J. (2015). Composition and potential origin of marine  
425 debris stranded in the Western Indian Ocean on remote Alphonse Island, Seychelles. Mar Pollut Bull, 96(1-  
426 2), 76-86. doi:10.1016/j.marpolbul.2015.05.042

427 Erftemeijer, P., Riegl, B., Hoeksema, B. and Todd, P. (2012). Environmental impacts of dredging and other  
428 sediment disturbances on corals: A review. Marine Pollution Bulletin, 64(9), pp.1737-1765.

429 Eriksen, M., Lebreton, L., Carson, H., Thiel, M., Moore, C., Borerro, J., Galgani, F., Ryan, P. and Reisser, J.  
 430 (2014). Plastic Pollution in the World's Oceans: More than 5 Trillion Plastic Pieces Weighing over 250,000  
 431 Tons Afloat at Sea. PLoS ONE, 9(12), p.e111913. doi:10.1371/journal.pone.0111913

432 Eriksson, C., Burton, H., Fitch, S., Schulz, M., and van den Hoff, J. (2013). Daily accumulation rates of marine  
 433 debris on sub-Antarctic island beaches. Mar Pollut Bull, 66(1-2), 199-208. doi:10.1016/j.marpol-  
 434 bul.2012.08.026

435 Eryasar, A. R., Ozbilgin, H., Gucu, A. C. and Sakinan, S. (2014). Marine debris in bottom trawl catches and  
 436 their effects on the selectivity grids in the north eastern Mediterranean. Mar Pollut Bull, 81(1), 80-84.  
 437 doi:10.1016/j.marpolbul.2014.02.017

438 European Commission (2015). Closing the loop - An EU action plan for the Circular Economy [pdf]. Available  
 439 from: [http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX: 52015DC0614](http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:52015DC0614) [Accessed 06/11/2016].

440 Fastelli, P., Blašković, A., Bernardi, G., Romeo, T., Čížmek, H., Andaloro, F., Russo, G., Guerranti, C. and  
 441 Renzi, M. (2016). Plastic litter in sediments from a marine area likely to become protected (Aeolian Archi-  
 442 pelago's islands, Tyrrhenian sea). Marine Pollution Bulletin, 113(1-2), pp.526-529. doi:10.1016/j.marpol-  
 443 bul.2016.08.054

444 Fernandino, G., Elliff, C., Silva, I., Brito, T. and Bittencourt, A. (2016). Plastic fragments as a major compo-  
 445 nent of marine litter: a case study in Salvador, Bahia, Brazil. Revista de Gestão Costeira Integrada, 16(3),  
 446 pp.281-287. doi:10.5894/rgci649

447 Fishing for litter Scotland (2015). Final Report 2011 - 2014. [online] Available at: [http://www.fishingforlit-  
 448 ter.org.uk/assets/file/Report%20FFL%202011%20-%202014.pdf](http://www.fishingforlitter.org.uk/assets/file/Report%20FFL%202011%20-%202014.pdf) [Accessed 3 Aug. 2017].

449 Fishing for litter Scotland (2017). Fishing for litter Scotland 2014 – 2017. [online] Available at:  
 450 <http://www.fishingforlitter.org.uk/assets/file/FFLS%202015.pdf> [Accessed 3 Aug. 2017].

451 Fishing for litter South West (2015). SW Update Dec 2015. [online] Available at: [http://www.fishingforlit-  
 452 ter.org.uk/assets/file/SW%20Update%20Dec%202015.pdf](http://www.fishingforlitter.org.uk/assets/file/SW%20Update%20Dec%202015.pdf) [Accessed 3 Aug. 2017].

453 Foekema, E. M., De Grujter, C., Mergia, M. T., van Franeker, J. A., Murk, A. J., and Koelmans, A. A. (2013).  
 454 Plastic in north sea fish. Environ Sci Technol, 47(15), 8818-8824. doi:10.1021/es400931b

455 Fukuoka, T., Yamane, M., Kinoshita, C., Narazaki, T., Marshall, G., Abernathy, K., Miyazaki, N. and Sato, K.  
 456 (2016). The feeding habit of sea turtles influences their reaction to artificial marine debris. Scientific Re-  
 457 ports, 6(1). doi:10.1038/srep28015

458 FUNDY NORTH Fisherman's Association (2016). Lost at Sea: A ghost gear retrieval manual. [online] Available  
 459 at: <http://www.fundynorth.org/wp-content/uploads/2016/03/Ghost-Gear-Retrieval-Manual2.pdf> [Ac-  
 460 cessed 3 Aug. 2017].

461 Gall, S. C. and Thompson, R. C. (2015). The impact of debris on marine life. Mar Pollut Bull, 92(1-2), 170-  
 462 179. doi:10.1016/j.marpolbul.2014.12.041

463 Gauquie, J., Devriese, L., Robbens, J., and De Witte, B. (2015). A qualitative screening and quantitative  
 464 measurement of organic contaminants on different types of marine plastic debris. Chemosphere, 138, 348-  
 465 356. doi:10.1016/j.chemosphere.2015.06.029

466 Ghost nets Australia (n.d.). Fact Sheets - GhostNets Australia. [online] Ghostnets.com.au. Available at:  
 467 <https://www.ghostnets.com.au/resources/fact-sheets/> [Accessed 3 Aug. 2017].

468 Gigault, J., Pedrono, B., Maxit, B., and Ter Halle, A. (2016). Marine plastic litter: the unanalyzed nano-frac-  
 469 tion. Environ. Sci.: Nano, 3(2), 346-350. doi:10.1039/c6en00008h

470 Gil, M. A., and Pfaller, J. B. (2016). Oceanic barnacles act as foundation species on plastic debris: implica-  
471 tions for marine dispersal. *Sci Rep*, 6, 19987. doi:10.1038/srep19987

472 Gilbert, J.M., Reichelt-Brushett, A. J., Bowling A.C., Christidis L. (2015). Plastic ingestion in marine and  
473 coastal bird species of Southeastern Australia. *Marine Ornithology*, 44, 21-26.

474 Goldstein, M. C., Titmus, A. J., and Ford, M. (2013). Scales of spatial heterogeneity of plastic marine debris  
475 in the northeast pacific ocean. *PLoS One*, 8(11), e80020. doi:10.1371/journal.pone.0080020

476 Goto, T. and Shibata, H. (2015). Changes in abundance and composition of anthropogenic marine debris on  
477 the continental slope off the Pacific coast of northern Japan, after the March 2011 Tohoku earthquake. *Mar*  
478 *Pollut Bull*, 95(1), 234-241. doi:10.1016/j.marpolbul.2015.04.011

479 Guerranti, C., Cannas, S., Scopetani, C., Fastelli, P., Cincinelli, A., and Renzi, M. (2017). Plastic litter in  
480 aquatic environments of Maremma Regional Park (Tyrrhenian Sea, Italy): Contribution by the Ombrone  
481 river and levels in marine sediments. *Mar Pollut Bull*, 117(1-2), 366-370. doi:10.1016/j.marpol-  
482 bul.2017.02.021

483 Hardesty, B. D., Lawson, T. J., van der Velde, T., Lansdell, M., and Wilcox, C. (2017). Estimating quantities  
484 and sources of marine debris at a continental scale. *Frontiers in Ecology and the Environment*, 15(1), 18-25.  
485 doi:10.1002/fee.1447

486 Hoarau, L., Ainley, L., Jean, C., and Ciccione, S. (2014). Ingestion and defecation of marine debris by logger-  
487 head sea turtles, *Caretta caretta*, from by-catches in the South-West Indian Ocean. *Mar Pollut Bull*, 84(1-2),  
488 90-96. doi:10.1016/j.marpolbul.2014.05.031

489 Iñiguez, M. E., Conesa, J. A. and Fullana, A. (2016). Marine debris occurrence and treatment: A review. *Re-*  
490 *newable and Sustainable Energy Reviews*, 64, 394-402. doi:10.1016/j.rser.2016.06.031

491 International Coastal Cleanup and Ocean Conservancy (2013). Working for clean beaches and clean water -  
492 2013 Report. [online] Available at: [http://coastalcleanup.nus.edu.sg/download/Final%20Re-](http://coastalcleanup.nus.edu.sg/download/Final%20Re-ports/2013_ICC_Report.pdf)  
493 [ports/2013\\_ICC\\_Report.pdf](http://coastalcleanup.nus.edu.sg/download/Final%20Re-ports/2013_ICC_Report.pdf) [Accessed 3 Aug. 2017].

494 International Coastal Cleanup and Ocean Conservancy (2014). Turning the tide on trash - 2014 Report.  
495 [online] Available at: [http://coastalcleanup.nus.edu.sg/download/Final%20Reports/2014\\_ICC\\_Report.pdf](http://coastalcleanup.nus.edu.sg/download/Final%20Reports/2014_ICC_Report.pdf)  
496 [Accessed 3 Aug. 2017].

497 International Coastal Cleanup and Ocean Conservancy (2015). Trash free seas every piece, every person... -  
498 2015 Report. [online] Available at: [http://coastalcleanup.nus.edu.sg/download/Final%20Re-](http://coastalcleanup.nus.edu.sg/download/Final%20Re-ports/2015_ICC_Report.pdf)  
499 [ports/2015\\_ICC\\_Report.pdf](http://coastalcleanup.nus.edu.sg/download/Final%20Re-ports/2015_ICC_Report.pdf) [Accessed 3 Aug. 2017].

500 International Coastal Cleanup and Ocean Conservancy (2016). 30th Anniversary International Coastal  
501 Cleanup - Annual Report. [online] Available at: [https://oceanconservancy.org/wp-content/up-](https://oceanconservancy.org/wp-content/uploads/2017/04/2016-data-release-1.pdf)  
502 [loads/2017/04/2016-data-release-1.pdf](https://oceanconservancy.org/wp-content/uploads/2017/04/2016-data-release-1.pdf) [Accessed 3 Aug. 2017].

503 Italian Institute of Marine Sciences (2016). Life Ghost - Technical deliverables. [online] Life-ghost.eu. Availa-  
504 ble at: [http://www.life-ghost.eu/index.php/en/downloads/technical-deliverables/send/4-technical-deliver-](http://www.life-ghost.eu/index.php/en/downloads/technical-deliverables/send/4-technical-deliver-ables/375-report-on-removal-activities)  
505 [ables/375-report-on-removal-activities](http://www.life-ghost.eu/index.php/en/downloads/technical-deliverables/send/4-technical-deliver-ables/375-report-on-removal-activities) [Accessed 3 Aug. 2017].

506 Jambeck, J. R., Geyer, R., Wilcox, C., Siegler, T. R., Perryman, M. and Andrady, A., Narayan, R., Law, K. L.  
507 (2015). Plastic waste inputs from land into the ocean. *Science*, 347(6223), 768-771.

508 Jambeck, J. and Johnsen, K. (2015). Citizen-Based Litter and Marine Debris Data Collection and Mapping.  
509 *Computing in Science & Engineering*, 17(4), pp.20-26.



510 Jang, Y. C., Lee, J., Hong, S., Lee, J. S., Shim, W. J., & Song, Y. K. (2014a). Sources of plastic marine debris on  
511 beaches of Korea: More from the ocean than the land. *Ocean Science Journal*, 49(2), 151-162.  
512 doi:10.1007/s12601-014-0015-8

513 Jang, Y., Lee, J., Hong, S., Mok, J., Kim, K., Lee, Y., Choi, H., Kang, H. and Lee, S. (2014b). Estimation of the  
514 annual flow and stock of marine debris in South Korea for management purposes. *Marine Pollution Bulletin*,  
515 86(1-2), pp.505-511. doi:10.1016/j.marpolbul.2014.06.021

516 Jantz, L. A., Morishige, C. L., Bruland, G. L., and Lepczyk, C. A. (2013). Ingestion of plastic marine debris by  
517 longnose lancetfish (*Alepisaurus ferox*) in the North Pacific Ocean. *Mar Pollut Bull*, 69(1-2), 97-104.  
518 doi:10.1016/j.marpolbul.2013.01.019

519 Jerdy, H., Werneck, M. R., da Silva, M. A., Ribeiro, R. B., Bianchi, M., Shimoda, E., and de Carvalho, E. C.  
520 (2017). Pathologies of the digestive system caused by marine debris in *Chelonia mydas*. *Mar Pollut Bull*,  
521 116(1-2), 192-195. doi:10.1016/j.marpolbul.2017.01.009

522 Jimenez, S., Domingo, A., Brazeiro, A., Defeo, O., and Phillips, R. A. (2015). Marine debris ingestion by alba-  
523 trosses in the southwest Atlantic Ocean. *Mar Pollut Bull*, 96(1-2), 149-154. doi:10.1016/j.marpol-  
524 bul.2015.05.034

525 Jones, M. M. (1995). Fishing debris in the Australian marine environment. *Mar. Pollut. Bull.*(30), 25–33.

526 Kadir, A.A., Hasni, A.F., and Sarani, N.A. (2015). Marine debris composition in Batupahat, Johor: A compari-  
527 son between Sungai Lurus and Minyakkbeku beaches. *ARPN Journal of Engineering and Applied Sciences*,  
528 10(15), 6553-6557.

529 Kalogerakis, N. K., S.; Yiantzi, E.; Perraki, T.; Kalogerakis, G.C.; Fodelianakis, S.;, and Fava, F. P., E. (2014).  
530 PCBs and PAHs on plastic pellets and microplastics collected on the coastline of the island of Crete in east-  
531 ern Mediterranean sea - Monitoring and fate. *Proceedings of the 37th AMOP Technical Seminar on Environ-  
532 mental Contamination and Response*.

533 Keswani, A., Oliver, D. M., Gutierrez, T. and Quilliam, R. S. (2016). Microbial hitchhikers on marine plastic  
534 debris: Human exposure risks at bathing waters and beach environments. *Mar Environ Res*, 118, 10-19.  
535 doi:10.1016/j.marenvres.2016.04.006

536 Kiessling, T., Gutow, L. and Thiel, M. (2015). Marine Litter as Habitat and Dispersal Vector. *Marine Anthro-  
537 pogenic Litter*. 141-181.

538 Kooi, M., Reisser, J., Slat, B., Ferrari, F., Schmid, M., Cunsolo, S., Brambini, R., Noble, K., Sirks, L., Linders, T.,  
539 Schoeneich-Argent, R. and Koelmans, A. (2016). The effect of particle properties on the depth profile of  
540 buoyant plastics in the ocean. *Scientific Reports*, 6(1). doi:10.1038/srep33882

541 Kühn, S., Bravo Rebolledo, E. L. and van Franeker, J. A. (2015). Deleterious Effects of Litter on Marine Life.  
542 *Marine Anthropogenic Litter*. 75-116.

543 Kwon, B. G., Koizumi, K., Chung, S. Y., Kodera, Y., Kim, J. O., and Saido, K. (2015). Global styrene oligomers  
544 monitoring as new chemical contamination from polystyrene plastic marine pollution. *J Hazard Mater*, 300,  
545 359-367. doi:10.1016/j.jhazmat.2015.07.039

546 Lavers, J. L., Hodgson, J. C., and Clarke, R. H. (2013). Prevalence and composition of marine debris in Brown  
547 Booby (*Sula leucogaster*) nests at Ashmore Reef. *Mar Pollut Bull*, 77(1-2), 320-324. doi:10.1016/j.marpol-  
548 bul.2013.09.026

549 Law, K. L. (2017). Plastics in the Marine Environment. *Ann Rev Mar Sci*, 9, 205-229. doi:10.1146/annurev-  
550 marine-010816-060409

551 Law, K. L., Moret-Ferguson, S. E., Goodwin, D. S., Zettler, E. R., Deforce, E., Kukulka, T., and Proskurowski, G.  
552 (2014). Distribution of surface plastic debris in the eastern Pacific Ocean from an 11-year data set. *Environ*  
553 *Sci Technol*, 48(9), 4732-4738. doi:10.1021/es4053076

554 Lawson, T. J., Wilcox, C., Johns, K., Dann, P., and Hardesty, B. D. (2015). Characteristics of marine debris that  
555 entangle Australian fur seals (*Arctocephalus pusillus doriferus*) in southern Australia. *Mar Pollut Bull*, 98(1-  
556 2), 354-357. doi:10.1016/j.marpolbul.2015.05.053

557 Lee, J., Lee, J., Jang, Y., Hong, S., Shim, W., Song, Y., Hong, S., Jang, M., Han, G., Kang, D. and Hong, S.  
558 (2015). Distribution and Size Relationships of Plastic Marine Debris on Beaches in South Korea. *Archives of*  
559 *Environmental Contamination and Toxicology*, 69(3), pp.288-298. doi:10.1007/s00244-015-0208-x

560 Leite, A. S., Santos, L. L., Costa, Y., and Hatje, V. (2014). Influence of proximity to an urban center in the pat-  
561 tern of contamination by marine debris. *Mar Pollut Bull*, 81(1), 242-247. doi:10.1016/j.marpol-  
562 bul.2014.01.032

563 Li, W. C., Tse, H. F. and Fok, L. (2016). Plastic waste in the marine environment: A review of sources, occur-  
564 rence and effects. *Sci Total Environ*, 566-567, 333-349. doi:10.1016/j.scitotenv.2016.05.084

565 Lieder, M. and Rashid, A. (2016). Towards circular economy implementation: a comprehensive review in  
566 context of manufacturing industry. *Journal of Cleaner Production*, 115, 36-51.

567 Lueiro, X. (2013). Sailing towards a 'Guardians of the Sea' horizon: CETMAR actions in marine litter with the  
568 fishing industry. [online] Available at: [http://www.marine-litter-conference-ber-](http://www.marine-litter-conference-berlin.info/userfiles/file/online/CETMAR%20actions%20in%20marine%20litter%20with%20the%20fish-ing%20industry_amoedo_lueiro.pdf)  
569 [lin.info/userfiles/file/online/CETMAR%20actions%20in%20marine%20litter%20with%20the%20fish-](http://www.marine-litter-conference-berlin.info/userfiles/file/online/CETMAR%20actions%20in%20marine%20litter%20with%20the%20fish-ing%20industry_amoedo_lueiro.pdf)  
570 [ing%20industry\\_amoedo\\_lueiro.pdf](http://www.marine-litter-conference-berlin.info/userfiles/file/online/CETMAR%20actions%20in%20marine%20litter%20with%20the%20fish-ing%20industry_amoedo_lueiro.pdf) [Accessed 3 Aug. 2017].

571 Marine Conservation Society (2016). Great British Beach Clean - 2016 Report. [online] Available at:  
572 [http://www.mcsuk.org/downloads/gbbc/2016/GBBC\\_2016\\_Report.pdf](http://www.mcsuk.org/downloads/gbbc/2016/GBBC_2016_Report.pdf) [Accessed 3 Aug. 2017].

573 Martin, J. M. (2013). Marine debris removal: one year of effort by the Georgia Sea Turtle-Center-Marine  
574 Debris Initiative. *Mar Pollut Bull*, 74(1), 165-169. doi:10.1016/j.marpolbul.2013.07.009

575 MARPOL (International Convention for the Prevention of Pollution from Ships), (1988). Annex V Prevention  
576 of Pollution by Garbage from Ships

577 Masó, M., Fortuño, J.-M., De Juan, S., and Demestre, M. (2016). Microfouling communities from pelagic and  
578 benthic marine plastic debris sampled across Mediterranean coastal waters. *Scientia Marina*, 80(S1), 117-  
579 127. doi:10.3989/scimar.04281.10A

580 McIlgorm, A., Campbell, H. F. and Rule, M. J. (2011). Understanding the economic benefits and costs of con-  
581 trolling marine debris in the APEC region. Singapore: APEC (MRC 02/2007).

582 Mobilik, J.-M., Ling, T.-Y., Husain, M.-L. B. and Hassan, R. (2015). Seasonal trends in abundance and compo-  
583 sition of marine debris in selected public beaches in Peninsular Malaysia. 1678, 020020.  
584 doi:10.1063/1.4931205

585 Naturschutzbund Deutschland (2015). Fishing for Litter: Müll fischen für saubere Meere - eine gemeinsame  
586 Initiative von Fischerei und Naturschutz. [online] Available at: [https://www.nabu.de/imperia/md/con-](https://www.nabu.de/imperia/md/content/nabude/naturschutz/meeresschutz/150312-nabu-infopapier-f4l.pdf)  
587 [tent/nabude/naturschutz/meeresschutz/150312-nabu-infopapier-f4l.pdf](https://www.nabu.de/imperia/md/content/nabude/naturschutz/meeresschutz/150312-nabu-infopapier-f4l.pdf) [Accessed 3 Aug. 2017].

588 Napper, I., Bakir, A., Rowland, S. and Thompson, R. (2015). Characterisation, quantity and sorptive proper-  
589 ties of microplastics extracted from cosmetics. *Marine Pollution Bulletin*, 99(1-2), pp.178-185.

590 National Fish and Wildlife Foundation (2016). Fishing for Energy. [online] Available at:  
591 [http://www.nfwf.org/fishingforenergy/Documents/ffe\\_factsheet\\_16-0329.pdf](http://www.nfwf.org/fishingforenergy/Documents/ffe_factsheet_16-0329.pdf) [Accessed 3 Aug. 2017].

NOAA Marine Debris Program (2015). Report on the impacts of “ghost fishing” via derelict fishing gear. Silver Spring, MD. 25 pp

Northwest Straits Foundation (2015). Shallow Water Legacy Derelict Net Removal from High Priority Areas of Puget Sound. [online] Available at: <http://www.nwstraitsfoundation.org/download/shallow-water-legacy-derelict-net-removal-from-high-priority-areas-of-puget-sound/> [Accessed 3 Aug. 2017].

Olive Ridley Project (2017). News from the field. [online] Available at: [http://oliveridleyproject.org/wp-content/uploads/2017/03/ORP-News\\_March-2017\\_3.pdf](http://oliveridleyproject.org/wp-content/uploads/2017/03/ORP-News_March-2017_3.pdf) [Accessed 3 Aug. 2017].

Pasternak, G., Zviely, D., Ribic, C. A., Ariel, A., and Spanier, E. (2017). Sources, composition and spatial distribution of marine debris along the Mediterranean coast of Israel. *Mar Pollut Bull*, 114(2), 1036-1045. doi:10.1016/j.marpolbul.2016.11.023

Pedrotti, M., Petit, S., Elineau, A., Bruzaud, S., Crebassa, J., Dumontet, B., Martí, E., Gorsky, G. and Cózar, A. (2016). Changes in the Floating Plastic Pollution of the Mediterranean Sea in Relation to the Distance to Land. *PLOS ONE*, 11(8), p.e0161581. doi:10.1371/journal.pone.0161581

Petry, M. V. and Benemann, V. R. (2017). Ingestion of marine debris by the White-chinned Petrel (*Procelaria aequinoctialis*): Is it increasing over time off southern Brazil? *Mar Pollut Bull*, 117(1-2), 131-135. doi:10.1016/j.marpolbul.2017.01.073

Polasek, L., Bering, J., Kim, H., Neitlich, P., Pister, B., Terwilliger, M., Nicolato, K., Turner, C. and Jones, T. (2017). Marine debris in five national parks in Alaska. *Marine Pollution Bulletin*, 117(1-2), pp.371-379. doi:10.1016/j.marpolbul.2017.01.085

Poli, C., Mesquita, D. O., Saska, C., and Mascarenhas, R. (2015). Plastic ingestion by sea turtles in Paraíba State, Northeast Brazil. *Iheringia. Série Zoologia*, 105(3), 265-270. doi:10.1590/1678-476620151053265270

Possatto, F. E., Spach, H. L., Cattani, A. P., Lamour, M. R., Santos, L. O., Cordeiro, N. M., and Broadhurst, M. K. (2015). Marine debris in a World Heritage Listed Brazilian estuary. *Mar Pollut Bull*, 91(2), 548-553. doi:10.1016/j.marpolbul.2014.09.032

Qari, R. S., M. (2015). Distribution and Abundance of Marine Debris Along the Coast of Karachi (Arabian Sea), Pakistan. *Pakistan Journal of Scientific and Industrial Research*, 58(2).

Rani, M., Shim, W. J., Han, G. M., Jang, M., Al-Odaini, N. A., Song, Y. K., and Hong, S. H. (2015). Qualitative Analysis of Additives in Plastic Marine Debris and Its New Products. *Arch Environ Contam Toxicol*, 69(3), 352-366. doi:10.1007/s00244-015-0224-x

Rani, M., Shim, W. J., Han, G. M., Jang, M., Song, Y. K., and Hong, S. H. (2017). Benzotriazole-type ultraviolet stabilizers and antioxidants in plastic marine debris and their new products. *Sci Total Environ*, 579, 745-754. doi:10.1016/j.scitotenv.2016.11.033

Reisser, J., Shaw, J., Hallegraeff, G., Proietti, M., Barnes, D., Thums, M., Wilcox, C., Hardesty, B. and Pattiaratchi, C. (2014). Millimeter-Sized Marine Plastics: A New Pelagic Habitat for Microorganisms and Invertebrates. *PLoS ONE*, 9(6), p.e100289. doi:10.1371/journal.pone.0100289

Reisser, J., Shaw, J., Wilcox, C., Hardesty, B., Proietti, M., Thums, M. and Pattiaratchi, C. (2013). Marine Plastic Pollution in Waters around Australia: Characteristics, Concentrations, and Pathways. *PLoS ONE*, 8(11), p.e80466. doi:10.1371/journal.pone.0080466

Reisser, J., Slat, B., Noble, K., du Plessis, K., Epp, M., Proietti, M., de Sonnevile, J., Becker, T. and Pattiaratchi, C. (2015). The vertical distribution of buoyant plastics at sea: an observational study in the North Atlantic Gyre. *Biogeosciences*, 12(4), pp.1249-1256.

633 Rochman, C. M., Lewison, R. L., Eriksen, M., Allen, H., Cook, A. M., and Teh, S. J. (2014). Polybrominated di-  
634 phenyl ethers (PBDEs) in fish tissue may be an indicator of plastic contamination in marine habitats. *Sci To-*  
635 *tal Environ*, 476-477, 622-633. doi:10.1016/j.scitotenv.2014.01.058

636 Romeo, T., Pietro, B., Peda, C., Consoli, P., Andaloro, F., and Fossi, M. C. (2015). First evidence of presence  
637 of plastic debris in stomach of large pelagic fish in the Mediterranean Sea. *Mar Pollut Bull*, 95(1), 358-361.  
638 doi:10.1016/j.marpolbul.2015.04.048

639 Ruiz-Oregon, L. F., Sarda, R., and Ramis-Pujol, J. (2016). Floating plastic debris in the Central and Western  
640 Mediterranean Sea. *Mar Environ Res*, 120, 136-144. doi:10.1016/j.marenvres.2016.08.001

641 Rummel, C. D., Loder, M. G., Fricke, N. F., Lang, T., Griebeler, E. M., Janke, M., and Gerdtts, G. (2016). Plastic  
642 ingestion by pelagic and demersal fish from the North Sea and Baltic Sea. *Mar Pollut Bull*, 102(1), 134-141.  
643 doi:10.1016/j.marpolbul.2015.11.043

644 Ryan, P. G., de Bruyn, P. J., and Bester, M. N. (2016). Regional differences in plastic ingestion among South-  
645 ern Ocean fur seals and albatrosses. *Mar Pollut Bull*, 104(1-2), 207-210. doi:10.1016/j.marpol-  
646 bul.2016.01.032

647 Santos, R. G., Andrades, R., Boldrini, M. A., and Martins, A. S. (2015). Debris ingestion by juvenile marine  
648 turtles: an underestimated problem. *Mar Pollut Bull*, 93(1-2), 37-43. doi:10.1016/j.marpolbul.2015.02.022

649 Santos, R. G., Andrades, R., Fardim, L. M., and Martins, A. S. (2016). Marine debris ingestion and Thayer's  
650 law - The importance of plastic color. *Environ Pollut*, 214, 585-588. doi:10.1016/j.envpol.2016.04.024

651 Sarafriz, J., Rajabizadeh, M., and Kamrani, E. (2016). The preliminary assessment of abundance and compo-  
652 sition of marine beach debris in the northern Persian Gulf, Bandar Abbas City, Iran. *Journal of the Marine*  
653 *Biological Association of the United Kingdom*, 96(01), 131-135. doi:10.1017/s0025315415002076

654 Sherrington, C., Darrah, C., Hann, S., Cole, G. and Corbin, M. (2016). Study to support the development of  
655 measures to combat a range of marine litter sources - Report for European Commission DG Environment  
656 [pdf]. Available from: [www.ec.europa.eu/ environment/marine/good-environmental-status/descriptor-](http://www.ec.europa.eu/environment/marine/good-environmental-status/descriptor-10/pdf/MSFD%20Measure%20to%20Combat%20Marine%20Litter.pdf)  
657 [10/pdf/MSFD%20Measure s%20to%20Combat%20Marine%20Litter.pdf](http://www.ec.europa.eu/environment/marine/good-environmental-status/descriptor-10/pdf/MSFD%20Measure%20to%20Combat%20Marine%20Litter.pdf) [Accessed 05/11/2016].

658 Smith, S. D., and Edgar, R. J. (2014). Documenting the density of subtidal marine debris across multiple ma-  
659 rine and coastal habitats. *PLoS One*, 9(4), e94593. doi:10.1371/journal.pone.0094593

660 Smith, S. D., Gillies, C. L., and Shortland-Jones, H. (2014). Patterns of marine debris distribution on the  
661 beaches of Rottnest Island, Western Australia. *Mar Pollut Bull*, 88(1-2), 188-193. doi:10.1016/j.marpol-  
662 bul.2014.09.007

663 Smith, S. D., and Markic, A. (2013). Estimates of marine debris accumulation on beaches are strongly af-  
664 fected by the temporal scale of sampling. *PLoS One*, 8(12), e83694. doi:10.1371/journal.pone.0083694

665 Swanson, R. L., Lwiza, K., Willig, K. and Morris, K. (2016). Superstorm Sandy marine debris wash-ups on  
666 Long Island - What happened to them? *Mar Pollut Bull*, 108(1-2), 215-231. doi:10.1016/j.marpol-  
667 bul.2016.04.029

668 Tanaka, K., Takada, H., Yamashita, R., Mizukawa, K., Fukuwaka, M. A., and Watanuki, Y. (2013). Accumula-  
669 tion of plastic-derived chemicals in tissues of seabirds ingesting marine plastics. *Mar Pollut Bull*, 69(1-2),  
670 219-222. doi:10.1016/j.marpolbul.2012.12.010

671 Tavares, D. C., da Costa, L. L., Rangel, D. F., de Moura, J. F., Zalmon, I. R., and Siciliano, S. (2016). Nests of  
672 the brown booby (*Sula leucogaster*) as a potential indicator of tropical ocean pollution by marine debris.  
673 *Ecological Indicators*, 70, 10-14. doi:10.1016/j.ecolind.2016.06.005

674 Taylor, M. L., Gwinnett, C., Robinson, L. F., and Woodall, L. C. (2016). Plastic microfibre ingestion by deep-  
675 sea organisms. *Sci Rep*, 6, 33997. doi:10.1038/srep33997

676 Tekman, M., Gutow, L., Macario, A., Haas, A., Walter, A. and Bergmann, M. (2017a). LITTERBASE: Online  
677 Portal for Marine Litter. [online] Litterbase.awi.de. Available at: <http://litterbase.awi.de/interaction> [Ac-  
678 cessed 10 Aug. 2017].

679 Tekman, M., Gutow, L., Macario, A., Haas, A., Walter, A. and Bergmann, M. (2017b). LITTERBASE: Online  
680 Portal for Marine Litter. [online] Litterbase.awi.de. Available at: [http://litterbase.awi.de/litter\\_graph](http://litterbase.awi.de/litter_graph) [Ac-  
681 cessed 16 November 2017].

682 Ter Halle, A., Ladirat, L., Gendre, X., Goudouneche, D., Pusineri, C., Routaboul, C., Tenailleau, C., Duployer,  
683 B. and Perez, E. (2016). Understanding the Fragmentation Pattern of Marine Plastic Debris. *Environmental*  
684 *Science & Technology*, 50(11), pp.5668-5675. doi:10.1021/acs.est.6b00594

685 Thiel, M., Hinojosa, I. A., Miranda, L., Pantoja, J. F., Rivadeneira, M. M., and Vasquez, N. (2013). Anthro-  
686 pogenic marine debris in the coastal environment: a multi-year comparison between coastal waters and local  
687 shores. *Mar Pollut Bull*, 71(1-2), 307-316. doi:10.1016/j.marpolbul.2013.01.005

688 Turner, A. (2016). Heavy metals, metalloids and other hazardous elements in marine plastic litter. *Mar Pol-*  
689 *lut Bull*, 111(1-2), 136-142. doi:10.1016/j.marpolbul.2016.07.020

690 Uchida, K., Hagita, R., Hayashi, T., and Tokai, T. (2016). Distribution of small plastic fragments floating in the  
691 western Pacific Ocean from 2000 to 2001. *Fisheries Science*, 82(6), 969-974. doi:10.1007/s12562-016-1028-  
692 2

693 Unger, B., Rebolledo, E., Deaville, R., Gröne, A., IJsseldijk, L., Leopold, M., Siebert, U., Spitz, J., Wohlsein, P.  
694 and Herr, H. (2016). Large amounts of marine debris found in sperm whales stranded along the North Sea  
695 coast in early 2016. *Marine Pollution Bulletin*, 112(1-2), pp.134-141. doi:10.1016/j.marpolbul.2016.08.027

696 United Nations General Assembly (2015). Resolution 70/1. Transforming our world: The 2030 Agenda for  
697 Sustainable Development. October 21. A/RES/70/1.

698 Van Cauwenberghe, L., Claessens, M., Vandegehuchte, M. B., Mees, J. and Janssen, C. R. (2013). Assess-  
699 ment of marine debris on the Belgian Continental Shelf. *Mar Pollut Bull*, 73(1), 161-169. doi:10.1016/j.mar-  
700 polbul.2013.05.026

701 Verlis, K. M., Campbell, M. L., and Wilson, S. P. (2013). Ingestion of marine debris plastic by the wedge-  
702 tailed shearwater *Ardenna pacifica* in the Great Barrier Reef, Australia. *Mar Pollut Bull*, 72(1), 244-249.  
703 doi:10.1016/j.marpolbul.2013.03.017

704 Verlis, K. M., Campbell, M. L., and Wilson, S. P. (2014). Marine debris is selected as nesting material by the  
705 brown booby (*Sula leucogaster*) within the Swain Reefs, Great Barrier Reef, Australia. *Mar Pollut Bull*, 87(1-  
706 2), 180-190. doi:10.1016/j.marpolbul.2014.07.060

707 Wedemeyer-Strombel, K. R., Balazs, G. H., Johnson, J. B., Peterson, T. D., Wicksten, M. K., and Plotkin, P. T.  
708 (2015). High frequency of occurrence of anthropogenic debris ingestion by sea turtles in the North Pacific  
709 Ocean. *Marine Biology*, 162(10), 2079-2091. doi:10.1007/s00227-015-2738-1

710 West, J. A., Hansen, G. I., Hanyuda, T., and Zuccarello, G. C. (2016). Flora of drift plastics: a new red algal  
711 genus, *Tsunami* *transpacifica* (Stylonematophyceae) from Japanese tsunami debris in the northeast Pacific  
712 Ocean. *Algae*, 31(4), 289-301. doi:10.4490/algae.2016.31.10.20

713 Wilson, S. P., and Verlis, K. M. (2017). The ugly face of tourism: Marine debris pollution linked to visitation  
714 in the southern Great Barrier Reef, Australia. *Mar Pollut Bull*, 117(1-2), 239-246. doi:10.1016/j.marpol-  
715 bul.2017.01.036

716 Woodall, L., Sanchez-Vidal, A., Canals, M., Paterson, G., Coppock, R., Sleight, V., Calafat, A., Rogers, A., Na-  
 717 ryanaswamy, B. and Thompson, R. (2014). The deep sea is a major sink for microplastic debris. *Royal Soci-*  
 718 *ety Open Science*, 1(4), pp.140317-140317. doi:10.1098/rsos.140317

719 WWF Poland (2015). Removal of derelict fishing gear, lost or discarded by fishermen in the Baltic Sea – Final  
 720 Report. [online] Available at: [http://d2ouvy59p0dg6k.cloudfront.net/downloads/wwf\\_poland\\_re-](http://d2ouvy59p0dg6k.cloudfront.net/downloads/wwf_poland_removal_of_derelict_fishing_gear_oct_2015_1.pdf)  
 721 [moval\\_of\\_derelict\\_fishing\\_gear\\_oct\\_2015\\_1.pdf](http://d2ouvy59p0dg6k.cloudfront.net/downloads/wwf_poland_removal_of_derelict_fishing_gear_oct_2015_1.pdf) [Accessed 3 Aug. 2017].

722 Yi, C. J. and Kannan, N. (2016). Solid Waste Transportation through Ocean Currents: Marine Debris Sightings  
 723 and their Waste Quantification at Port Dickson Beaches, Peninsular Malaysia. *Environment Asia*, 9(2), 39-  
 724 47.

725 Zhao, S., Zhu, L., and Li, D. (2015). Characterization of small plastic debris on tourism beaches around the  
 726 South China Sea. *Regional Studies in Marine Science*, 1, 55-62. doi:10.1016/j.rsma.2015.04.001

727 Zhou, C., Liu, X., Wang, Z., Yang, T., Shi, L., Wang, L., You, S., Li, M. and Zhang, C. (2016). Assessment of ma-  
 728 rine debris in beaches or seawaters around the China Seas and coastal provinces. *Waste Management*, 48,  
 729 pp.652-660. doi:10.1016/j.wasman.2015.11.010

Nr	Removed quantity		Ocean compartment	Collection method	Location	Removal period	Waste treatment	Collector / Reference
	Value	Unit						
1	10404	kg	Coastline	Manually	Alaska (USA)	2015	-	(Polasek et al., 2017)
2	422	kg	Seafloor	Bottom trawling	Pacific Ocean	2003-2011	-	(Goto and Shibata, 2015)
3	244	kg	Coastline	Manually	Caribbean Sea	2015	-	(De Scisciolo et al., 2016)
4	231	kg	Coastline	Manually	Malaysia	-	-	(Mobilik et al., 2015)
5	227	kg	Seafloor	Bottom trawling	Mediterranean Sea	2010-2012	-	(Eryasar et al., 2014)
6	181	kg	Coastline	Manually	Brazil	2011-2012	-	(Da Silva et al., 2015a)
7	167	kg	Coastline	Manually	Malaysia	2014	-	(Yi and Kannan, 2016)
8	142	kg	Coastline	Manually	Seychelles	2013	-	(Duhec et al., 2015)
9	130	kg	Coastline	Manually	India	2015	-	(Arun Kumar et al., 2016)
10	74	kg	Coastline	Manually	Belgium	2010-2011	-	(Van Cauwenberghe et al., 2013)
	<1	kg	Sea surface	Surface trawling				
	<1	kg	Seafloor	Bottom trawling				
11	39	kg	Biota	Necropsy	North Sea	2016	-	(Unger et al., 2016)
12	30	kg	Sea surface	Retention boom	Hawaii (USA)	2011-2012	-	(Carson et al., 2013)
13	30	kg	Coastline	Manually	Brazil	2013	-	(Da Silva et al., 2016)
14	19	kg	Coastline	Manually	Malaysia	-	-	(Kadir, Hasni and Sarani, 2015)
15	16	kg	Coastline	Manually	Brazil	-	-	(Leite et al., 2014)
16	13	kg	Coastline	Manually	Morocco	2015	-	(Alshawafi et al., 2017)
17	12	kg	Coastline	Manually	Pakistan	2012	-	(Qari, 2015)
18	12	kg	Coastline	Manually	South Korea	2013	-	(Jang et al., 2014a)
19	4	kg	Coastline	Manually	Caribbean Sea	2011	-	(Debrot et al., 2013)
20	1	kg	Coastline	Manually	United Kingdom	2015	-	(Turner, 2016)
21	<1	kg	Biota	Necropsy	Brazil	2009-2013	-	(Santos et al., 2015)
22	<1	kg	Biota	Necropsy	Pacific Ocean	1993-2011	-	(Wedemeyer-Strombel et al., 2015)
23	<1	kg	Biota	Necropsy	Brazil	2009-2010	-	(Poli et al., 2015)
24	<1	kg	Biota	Necropsy	Mediterranean Sea	2005-2015	-	(Casale et al., 2016)
25	<1	kg	Biota	Necropsy	Mediterranean Sea	2010-2011	-	(Campani et al., 2013)
26	<1	kg	Biota	Necropsy	Pacific Ocean	2010-2011	-	(Jantz et al., 2013)
27	<1	kg	Sea surface	Surface trawling	Atlantic Ocean	2015	-	(Kooi et al., 2016)
28	<1	kg	Sea surface	Surface trawling	Atlantic Ocean	2014	-	(Ter Halle et al., 2016)
29	<1	kg	Biota	Necropsy	Washington (USA), British Columbia (Canada)	2001-2011	-	(Avery-Gomm et al., 2013)
30	<1	kg	Biota	Necropsy	Atlantic Ocean	2011-2012	-	(Bond et al., 2013)



31	179904	items	Coastline	Flotation	Hong Kong	2014-2015	-	(Cheung, Cheung and Fok, 2016)
32	69122	items	Coastline	Manually	Israel	2012-2015	-	(Pasternak et al., 2017)
33	46728	items	Coastline	Manually	Mediterranean Sea	2011-2013	-	(Smith and Markic, 2013)
34	32414	items	Sea surface	Surface trawling	Pacific Ocean	2009-2010	-	(Goldstein, Titmus and Ford, 2013)
35	17089	items	Coastline	Manually	Brazil	2012-2013	-	(Fernandino et al., 2016)
36	8055	items	Coastline	Manually	Chile	2002-2005	-	(Thiel et al., 2013)
37	6527	items	Coastline	Manually	Georgia (USA)	2012-2013	-	(Martin, 2013)
38	6389	items	Coastline	Manually	Sub-Antarctic island	2000-2001	-	(Eriksson et al., 2013)
39	6030	items	Sediments	Diving	Mediterranean Sea	2015	-	(Blaskovic et al., 2017)
40	4618	items	Coastline	Manually	Australia	2011-2013	-	(Hardesty et al., 2017)
41	4520	items	Coastline	Manually	Australia	2012	-	(Smith, Gillies, and Shortland-Jones, 2014)
42	3901	items	Sea surface	Surface trawling	Mediterranean Sea	2013	-	(Cozar et al., 2015)
43	3070	items	Sea surface	Surface trawling	International	2010-2011	-	(Cozar et al., 2014)
44	2986	items	Seafloor	Diving	Australia	2005-	-	(Smith and Edgar, 2014)
45	2673	items	Sea surface	Surface trawling	Mediterranean Sea	2012-2013	-	(De Lucia et al., 2014)
46	1992	items	Coastline	Manually	Brazil	-	-	(Santos et al., 2016)
	656	items	Biota	Necropsy				
47	1315	items	Biota	Necropsy	Indian Ocean	2007-2013	-	(Hoarau et al., 2014)
48	839	Items	Sea surface	Surface trawling	Australia	2011-2012	-	(Reisser et al., 2013)
49	587	items	Coastline	Flotation	China	2014	-	(Zhao, Zhu, and Li, 2015)
50	553	items	Biota	Necropsy	Brazil	1990-2014	-	(Petry and Benemann, 2017)
51	493	items	Coastline	Manually	Brazil	2016	-	(Tavares et al., 2016)
52	291	items	Seafloor	Bottom trawling	Atlantic Ocean	-	-	(Possatto et al., 2015)
53	162	items	Sea surface	Surface trawling	Pacific Ocean	2000-2001	-	(Uchida et al., 2016)
54	138	items	Biota	Free (entanglement)	Australia	1997-2012	-	(Lawson et al., 2015)
55	84	items	Coastline	Manually	Ashmore Reef	2013	-	(Lavers, Hodgson and Clarke, 2013)
56	68	items	Sea surface	Surface trawling	Australia	-	-	(Reisser et al., 2014)
57	50	items	Biota	Necropsy	Pacific Ocean	-	-	(Gilbert et al., 2015)
58	26	items	Sea surface	Manually	Mediterranean Sea	2009	-	(Masó et al., 2016)
	16	items	Seafloor	Bottom trawling				
59	39	items	Biota	Necropsy	Uruguay	2005-2013	-	(Jimenez et al., 2015)

60	37	items	Biota	Necropsy	Mediterranean Sea	2010	-	(Anastasopoulou et al., 2013)
61	29	items	Biota	Necropsy	Mediterranean Sea	2012-2013	-	(Romeo et al., 2015)
62	29	items	Coastline	Manually	South Korea	2014	-	(Rani et al., 2017)
63	19	items	Biota	Necropsy	Atlantic Ocean	-	-	(Di Benedetto and Ramos, 2014)
64	19	items	Coastline	Manually	South Korea	2014	-	(Rani et al., 2015)
65	17	items	Biota	Necropsy	Philippines	2015	-	(Abreo et al., 2016)
66	1360	dm <sup>3</sup>	Coastline	Manually	Caribbean Sea	2014	-	(Bennett-Martin, Visaggi, and Hawthorne, 2015)
67	-	-	Coastline	Manually	International	-	-	(Kwon et al., 2015)
	170	dm <sup>3</sup>	Sea surface	Manually				
68	56	dm <sup>3</sup>	Sediments	Manually	Mediterranean Sea	2015-2016	-	(Guerranti et al., 2017)
69	-	-	Sea surface	Surface trawling	Caribbean Sea	2013-2014	-	(Correa-Herrera et al., 2017)
70	-	-	Biota	Necropsy	Brazil	2010-2013	-	(Jerdy et al., 2017)
71	-	-	Coastline	Manually	Australia	2012	-	(Wilson and Verlis, 2017)
72	-	-	Coastline	Manually	Iran	2005	-	(Sarafraz, Rajabizadeh and Kamrani, 2016)
73	-	-	Biota	Necropsy	Ireland	2012-2016	-	(Acampora et al., 2016)
74	-	-	Biota	Necropsy	Mediterranean Sea	2012	-	(Battaglia et al., 2015)
75	-	-	Sediment	Diving	Mediterranean Sea	-	-	(Fastelli et al., 2016)
76	-	-	Biota	Necropsy	Japan	2012-2015	-	(Fukuoka et al., 2016)
77	-	-	Sea surface	Surface trawling	Atlantic Ocean	2014	-	(Gigault et al., 2016)
78	-	-	Sea surface	Manually	Pacific Ocean	2012	-	(Gil and Pfaller, 2016)
79	-	-	Sea surface	Surface trawling	Mediterranean Sea	2013	-	(Pedrotti et al., 2016)
80	-	-	Sea surface	Surface trawling	Mediterranean Sea	2011-2013	-	(Ruiz-Oregon, Sarda, and Ramis-Pujol, 2016)
81	-	-	Biota	Necropsy	North Sea, Baltic Sea	2013	-	(Rummel et al., 2016)
82	-	-	Biota	Necropsy	South Africa	1989 - 2014	-	(Ryan, de Bruyn, and Bester, 2016)
83	-	-	Sediment	ROV	Pacific Ocean	2011 - 2013	-	(Taylor et al., 2016)
84	-	-	Coastline	Manually	Washington (USA), Oregon (USA)	2012-2016	-	(West et al., 2016)
85	-	-	Coastline	Manually	China	2007-2013	-	(Zhou et al., 2016)
	-	-	Sea surface	Surface trawling				
86	-	-	Coastline	Manually	Hawaii (USA)	1990-2012	-	(Agustin et al., 2015)
87	-	-	Biota	Necropsy	Brazil	2008-2009	-	(Da Silva et al., 2015b)
88	-	-	Coastline	Manually	Washington (USA), British Columbia (Canada)	2008 - 2012	-	(Davis and Murphy, 2015)
	-	-	Sea surface	Surface trawling				

89	-	-	Biota	Necropsy	Brazil	2011	-	(De Carvalho et al., 2015)
90	-	-	Coastline	Manually	North Sea	2014	-	(De Tender et al., 2015)
			Water column	Niskin bottles				
			Seafloor	Bottom trawling				
			Sediment	Grabs				
91	-	-	Sea surface	Surface trawling	Atlantic Ocean	2014	-	(Reisser et al., 2015)
92	-	-	Coastline	Manually	Belgium	-	-	(Gauquie et al., 2015)
93	-	-	Coastline	Manually	South Korea	2013-2014	-	(Lee et al., 2015)
94	-	-	Coastline	Manually	Australia	2012-2013	-	(Verlis, Campbell and Wilson, 2014)
95	-	-	Biota	Necropsy	Brazil	-	-	(Di Benedetto and Awabdi, 2014)
96	-	-	Sea surface	Surface trawling	International		-	(Eriksen et al., 2014)
97	-	-	Sea surface	Surface trawling	Pacific Ocean	2001-2012	-	(Law et al., 2014)
98	-	-	Sea surface	Surface trawling	Atlantic Ocean	2010	-	(Rochman et al., 2014)
	-	-	Biota	Necropsy				
99	-	-	Sediment	ROV	International	2001-2012	-	(Woodall et al., 2014)
100	-	-	coastline	Manually	Mediterranean Sea	2013-2014	-	(Kalogerakis and Fava, 2014)
101	-	-	Biota	Necropsy	North Sea	2010-2011	-	(Foekema et al., 2013)
102	-	-	Biota	Necropsy	Pacific Ocean		-	(Tanaka et al., 2013)
103	-	-	Biota	Necropsy	Australia	2012	-	(Verlis, Campbell and Wilson, 2013)

732 **Appendix 2 – Extracted data from non-scientific collections**

Nr	Removed quantity		Ocean compartment	Collection method	Location	Removal period	Waste treatment	Collector / Reference
	Value	Unit						
1	176940	tonnes	Coastline	-	New York (USA)	2012	Energy recovery, landfill	Local authorities (Swanson et al. 2016)
2	27297	tonnes	Coastline	-	South Korea	annually	-	Local authorities (Jang et al. 2014b)
	3946	tonnes	Sea surface	-				
	11370	tonnes	Seafloor	-				
3	10285	tonnes	Coastline	-	South Korea	2000-2006	-	Korean Coastal cleanup campaigns (NOAA Marine Debris Program 2015)
4	8193	tonnes	Coastline	Manually	International	2015	-	International Coastal Cleanup and Ocean Conservancy (2016)
5	7342	tonnes	Coastline	Manually	International	2014	-	International Coastal Cleanup and Ocean Conservancy (2015)
6	5592	tonnes	Coastline	Manually	International	2013	-	International Coastal Cleanup and Ocean Conservancy (2014)
7	4603	tonnes	Coastline	Manually	International	2012	-	International Coastal Cleanup and Ocean Conservancy (2013)
8	1300	tonnes	Sea surface	Retention boom	South Korea	annually	Recycling, energy recovery, landfill	Local authorities (Iñiguez, Conesa, and Fullana 2016)
	350	tonnes	Seafloor	Crane excavation				
9	908	tonnes	Seafloor	Bottom trawling	Atlantic Ocean, North Sea	2005-2015	Landfill	Fishing for litter Scotland (2015, 2017)
10	820	tonnes	Seafloor	-	Hawaii (USA)	1996-2014	-	Pacific Islands Fisheries Science Center (NOAA Marine Debris Program 2015)
11	268	tonnes	Seafloor	Bottom trawling	Baltic Sea	2015	-	WWF Poland (2015)
12	250	tonnes	Seafloor	-	USA	2008	Recycling of metal, energy recovery, landfill	National Fish and Wildlife Foundation (2016)
13	104	tonnes	Seafloor	Diving	International	2014-2015	-	Project AWARE (International Coastal Cleanup and Ocean Conservancy, 2015; 2016)
14	67	tonnes	Seafloor	Bottom trawling	United Kingdom	2014-2015	-	Fishing for litter South West (2015)
15	60+	tonnes	Seafloor	-	California (USA)	2006-2012	-	CA lost fishing gear recovery project (NOAA Marine Debris Program 2015)

16	37+	tonnes	Seafloor	ROV + Diving	Washington (USA)	2002-2016	Recycling of lead line, landfill	Northwest Straits Foundation (2015)
17	34	tonnes	Seafloor	Bottom trawling	Spain	2009-2010	-	CETMAR (Lueiro, 2013)
18	20	tonnes	Seafloor	-	North Sea, Mediterranean Sea	2000-2006	-	Healthy Seas Initiative (NOAA Marine Debris Program 2015)
19	12	tonnes	Coastline	-	Hawaii (USA)	2014	-	US marine park Papahānaumokuākea (Cressey, 2016)
20	6+	tonnes	Sea surface	Surface trawling	Australia	2004-2014	Recycling into art, landfill	Ghost nets Australia (n.d.)
21	2+	tonnes	Seafloor	Bottom trawling	North Sea, Baltic Sea	2011-2014	-	Naturschutzbund Deutschland (2015)
22	2+	tonnes	Seafloor	Bottom trawling	New Brunswick (Canada)	2008-2015	Reuse fishing gear, recycling of rope, landfill	FUNDY NORTH Fisherman's Association (2016)
23	1	tonnes	Seafloor	-	Pakistan, Maldives	<2016	Recycling into art	Olive Ridley Project (2017)
24	1	tonnes	Seafloor	Diving	Mediterranean Sea	2014-2016	-	Italian Institute of Marine Sciences (2016)
25	400,968	items	Coastline	Manually	International	2011-2014	-	Marine debris tracker (Jambeck and Johnsen 2015)
26	268384	items	Coastline	Manually	United Kingdom	2016	-	Marine Conservation Society (2016)
27	33297	items	Seafloor	-	Virginia (USA)	2008-2013	-	CCRM VIMS (NOAA Marine Debris Program 2015)
28	12,000+	items	Seafloor	-	Australia	2004-2012	-	GhostNets Australia (NOAA Marine Debris Program 2015)
29	5,600	items	Seafloor	-	Maine (USA)	2000-2006	-	Gear Grab.org (NOAA Marine Debris Program 2015)